

# UNCLASSIFIED

AD NUMBER
AD832784
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAR 1968. Other requests shall be referred to Commanding Officer, Fort Detrick, Attn: Technical Releases Branch, TID, Fort Detrick, Frederick, MD 21701.
AUTHORITY
BDRL D/A ltr, 29 Sep 1971

THIS PAGE IS UNCLASSIFIED


AD832784

AD

TECHNICAL MANUSCRIPT 452

GROWTH OF RIFT VALLEY FEVER VIRUS  
IN HUMAN DIPLOID (WI-38) CELLS

Roger W. Johnson  
Michael D. Orlando  
William C. Patrick, III



MARCH 1968

STATEMENT #2 UNCLASSIFIED

This document is subject to special export controls and each  
transmittal to foreign governments or foreign nationals may be  
made only with prior approval of -----


DEPARTMENT OF THE ARMY

Fort Detrick

Frederick, Maryland

*Att: Tech. Release Br. - Tech. Info. Div.*

# DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST  
QUALITY AVAILABLE. THE COPY  
FURNISHED TO DTIC CONTAINED  
A SIGNIFICANT NUMBER OF  
PAGES WHICH DO NOT  
REPRODUCE LEGIBLY.**

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DISTRIBUTION TO	
CPSTI	WHITE SECTION <input checked="" type="checkbox"/>
DDC	DDC SECTION <input checked="" type="checkbox"/>
UNANNOUNCED	<input type="checkbox"/>
JUSTIFICATION	
BY	
DISTRIBUTION/AVAILABILITY CODES	
DDC	AVAIL. and/or SPECIAL
2	

DEPARTMENT OF THE ARMY  
Fort Detrick  
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 452

GROWTH OF RIFT VALLEY FEVER VIRUS IN HUMAN DIPLOID (WI-38) CELLS

Roger W. Johnson  
Michael D. Orlando  
William C. Patrick, III

Product Development Division  
AGENT DEVELOPMENT AND ENGINEERING LABORATORY

Project 1B533001D426

March 1968

In conducting the research described in this report, the investigators adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences-National Research Council.

#### ACKNOWLEDGMENT

We acknowledge the technical assistance of Mr. Jerry Keller.

#### ABSTRACT

The growth of Rift Valley fever virus in human diploid (WI-38) cells was studied. Peak titers of  $1 \times 10^{9.0}$  and  $1 \times 10^{9.4}$  mouse intracerebral median lethal doses per milliliter were obtained in monolayer culture and suspensions, respectively.

# GROWTH OF RIFT VALLEY FEVER VIRUS IN HUMAN DIPLOID (WI-38) CELLS

The growth of Rift Valley fever (RVF) virus in established cell lines<sup>1,2</sup> and in primary cells<sup>3,4</sup> has been described. The present report describes the growth of RVF virus in the human diploid strain WI-38.<sup>5</sup>

We used strain WI-38 cells in the 22nd passage,\* either as monolayer cultures in 32-oz glass prescription bottles or in suspension in Eagle's basal medium<sup>6</sup> supplemented with 10% calf serum.

The Van Wyk pantropic strain of RVF virus,<sup>7</sup> passed once in tissue culture from infected lamb serum and stored at -80 C, was used.

Washed monolayer cultures were infected with enough RVF virus in 10-ml amounts to give an input multiplicity of 15 MICLD<sub>50</sub> per cell. After 1 hour adsorption at 37 C, the inoculum was poured off. The cultures were then washed three times, and 80 ml were added of either Eagle's basal medium or medium 199,<sup>8</sup> both supplemented with 10% calf serum and 50 µg kanamycin\*\*/ml. The cultures were incubated at 37 C and assayed periodically for virus by mouse intracerebral inoculation.

Suspensions of WI-38 cells, adjusted to a density of about  $6 \times 10^5$  cells/ml in 500-ml volumes, were placed in 1-liter cylinders fitted with New Brunswick fermentor heads (impeller housing, vent and sample line). These suspensions were infected after 10 minutes' agitation (impeller speed 100 to 125 rpm) with enough virus to yield an input multiplicity of about 60 MICLD<sub>50</sub> per cell. The cultures were sampled periodically for cell count and virus assay.

In monolayer culture (Fig. 1), after an initial period of about 6 hours characterized by little increase in viral titer, the titer increased rapidly, reaching a peak of about  $1 \times 10^{9.0}$  MICLD<sub>50</sub>/ml by 30 hours. The titer then remained constant through the 72nd hour, the last time the cultures were assayed. Neither Eagle's basal medium nor medium 199 had any appreciable effect on virus growth in the WI-38 cells.

In suspensions of WI-38 cells (Fig. 2), after a lag of about 6 hours, viral titer increased to a peak of about  $1 \times 10^{9.4}$  by 27 hours. The titer remained constant until the 76th hour, after which it decreased. The viable cell counts decreased over the course of the study, reaching zero by 94 hours postinfection.

\* Received from Flow Laboratories, Inc., Rockville, Md.

\*\* Kantrex, Bristol Laboratory, Syracuse, N.Y.




Figure 1. Growth of Rift Valley Fever Virus in Human Diploid Cell Monolayer Culture. ●—● = Growth in Eagle's basal medium; ○—○ = growth in medium 199; both media were supplemented with 10% calf serum and 50  $\mu$ g kanamycin/ml.






Figure 2. Growth of Rift Valley Fever Virus in Suspended Human Diploid Cells.  
 ●—● = Virus titer; ○--○ = viable cell count.

The growth curves of RVF virus in monolayer cultures and suspensions of WI-38 cells were essentially the same as those for RVF virus in monolayer and suspension cultures of mouse fibroblast-like (MFL) cells.<sup>3</sup> The titer of the virus in the suspended WI-38 cells remained high longer than it did in suspensions of MFL cells<sup>3</sup> (50 hours versus 15 hours from peak, respectively). This could be caused in part by the difference in the media used in the two studies (Eagle's basal medium in the present study, medium 199 in the previous) and by a difference in impeller speeds (100 to 125 rpm in the present study compared with about 250 to 300 rpm in the previous study).

The viability of the infected WI-38 cells in suspension followed a course similar to that of the infected MFL cells in suspension.

Should serially transferred cell strains ever be approved for use in vaccine production, the methods of growing virus described in this report should prove easier and more economical and provide a more standard cell population than do the primary cells currently used for RVF virus vaccine production.<sup>4</sup>

LITERATURE CITED

1. Easterday, B.C.; Murphy, L.C. 1963. The growth of Rift Valley fever virus in cultures of established lines of cells. *Cornell Vet.* 53:3-11.
2. Johnson, R.W.; Orlando, M.D. 1968. Growth of Rift Valley fever virus in tissue culture. *Amer. J. Vet. Res.* 29:463-472.
3. Randall, K.; Gibbs, C.J., Jr.; Aulisio, C.G.; Binn, L.N.; Harrison, V.R. 1962. The development of a formalin-killed Rift Valley fever virus vaccine for use in man. *J. Immunol.* 89:660-671.
4. Binn, L.N.; Randall, R.; Harrison, V.R.; Gibbs, C.J., Jr.; Aulisio, C.G. 1963. Immunization against Rift Valley fever: The development of vaccines from nonprimate cell cultures and chick embryos. *Amer. J. Hyg.* 77:160-168.
5. Hayflick, L.; Moorehead, P.S. 1961. The serial cultivation of human diploid cell strains. *Exp. Cell Res.* 25:585-621.
6. Eagle, H. 1955. Nutrition needs of mammalian cells in tissue culture. *Science* 122:501-504.
7. Kaschula, V.R. 1953. The propagation and modification of strains of Rift Valley fever viruses in embryonated eggs and their use as immunizing agents for domestic ruminants. Doctor of Veterinarian Science Thesis, University of Pretoria, Onderstepoort, S. Africa.
8. Morgan, J.F.; Morton, H.J.; Parker, R.C. 1950. Nutrition of animal cells in tissue culture: I. Initial studies on a synthetic medium. *Proc. Soc. Exp. Biol. Med.* 73:1-8.

Unclassified

11

Security Classification

DOCUMENT CONTROL DATA - R & D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Department of the Army Fort Detrick, Frederick, Maryland, 21701		Unclassified
		2b. GROUP
3. REPORT TITLE		
GROWTH OF RIFT VALLEY FEVER VIRUS IN HUMAN DIPLOID (WI-38) CELLS		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name)		
Roger W. Johnson Michael D. Orlando William C. Patrick, III		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
March 1968	11	8
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S)
a. PROJECT NO. 1B533001D426		Technical Manuscript 452
c.		8b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)
d.		
9. DISTRIBUTION STATEMENT		
Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
		Department of the Army Fort Detrick, Frederick, Maryland, 21701
13. ABSTRACT		
The growth of Rift Valley fever virus in human diploid (WI-38) cells was studied. Peak titers of $1 \times 10^{5.0}$ and $1 \times 10^{9.4}$ mouse intracerebral median lethal doses per milliliter were obtained in monolayer culture and suspensions, respectively.		
14. Key Words		
*Rift Valley fever virus *Diploid *Humans Tissue culture		

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified

Security Classification